# FIFTH SEMESTER B.Sc. DEGREE EXAMINATION, NOVEMBER 2019

(CUCBCSS-UG)

Mathematics

MAT 5B 05-VECTOR CALCULUS

Time : Three Hours

Maximum: 120 Marks

### Part A

Answer all the twelve questions. Each question carries 1 mark.

- 1. Evaluate  $\lim_{(x,y)\to(1,3)} \frac{x+1}{4-y}$ .
- 2. Find the domain and range of  $z = \sqrt{1 x^2 y^2}$ .
- 3. Find the gradient of  $\phi(x, y, z) = x^2 + y^2 + z^2$ .
- 4. Compute the divergence of  $\vec{f} = xy \vec{i} + yz \vec{j} + xz \vec{h}$ .
- 5. Define directional derivative of a function.
- 6. What do you mean by a conservative vector field?
- 7. Give a very brief discription of linearization of a function of two variables.
- 8. Find du if  $u = e^{x^2 + y^2 + z^2}$ .
- 9. Fill in the blanks : If  $\hat{f}$  and  $\hat{g}$  are differentiable vector point functions, then

$$\nabla \cdot (\vec{f} \times \vec{g}) = \dots$$

10. State the tangential form of Green's theorem in the plane.

Turn over

- 11. Fill in the blanks: If  $\vec{a}$  is a constant vector and  $\vec{r} = x \vec{i} + y \vec{j} + z \vec{h}$ , then  $\nabla \times (\vec{a} \times \vec{r}) = \dots$
- 12. State Stokes theorem mentioning all the assumptions involved in it explicitly.

 $(12 \times 1 = 12 \text{ marks})$ 

## Part B

Answer any ten questions. Each question carries 4 marks.

- 13. Find the vector normal to the surface  $\phi(x, y, z) = x^2y 2y^2z^3$  at (1, -1, 2).
- 14. Evaluate  $\lim_{(x,y)\to(0,0)} \frac{x^2-xy}{\sqrt{x}-\sqrt{y}}.$
- 15. If  $x^2 + y^2 + z^2 + ye^x$   $z + z \cos y = 0$  then, find  $\frac{\partial z}{\partial x}$  and  $\frac{\partial z}{\partial y}$  at the origin.
- 16. Prove that  $\nabla (r^n) = nr^{n-2} \vec{r}$ .
- 17. Find the total derivative of u = xy + z with respect to t if  $x = \cos t$ ,  $y = \sin t$  and z = t.
- 18. Compute the average value of the function  $f(x, y) = x \cos(xy)$  over the rectangular region  $0 \le x \le \pi, 0 \le y \le 1$ .
- 19. Linearize the function  $f(x, y) = \sin(\pi xy^2)$  at (1, 1).
- 20. Find the directional derivative of  $f(x, y) = xe^y + \cos(xy)$  at (2,0) in the direction of 3i 4j.
- 21. Find the velocity and acceleration vectors of  $r(t) = (3\cos t)i + (3\sin t)j + t^2k$ .
- 22. Find the flow of  $\vec{r} = x \vec{i} + y \vec{j} + z \vec{k}$  along the portion of the circular helix  $x = \cos t$ ,  $y = \sin t$ , z = t;  $0 \le t \pi/2$ .

- 23. Test whether the vector  $\vec{f} = (e^x \cos y + yz)\vec{i} + (xz e^x \sin y)\vec{j} + (xy + z)\vec{k}$  is conservative or not.
- 24. If the sides and angles in a traingle vary in such a way that its circum-radius R remains a constant, then show that  $\frac{da}{\cos A} + \frac{db}{\cos B} + \frac{dc}{\cos C} = 0$ .
- 25. Verify whether the differential ydx + xdy + 4dz is exact or not.
- 26. Show that  $\vec{f} \times \vec{g}$  is solenoidal if  $\vec{f}$  and  $\vec{g}$  are irrotational.

 $(10 \times 4 = 40 \text{ marks})$ 

# Part C

Answer any six questions. Each question carries 7 marks.

- 27. Evaluate  $\int_0^\pi \int_y^\pi \frac{\sin x}{x} dy dx$ .
- 28. If  $\vec{f}$  is a differentiable vector function of t, differentiable at least 3 times, prove that  $\frac{d}{dt} \left[ \vec{f}, \vec{f}', \vec{f}'' \right] = \left[ \vec{f}', \vec{f}'', \vec{f}''' \right].$
- 29. Find the work done by the force field  $\vec{f} = z\vec{i} + x\vec{j} + y\vec{k}$  along the boundary of the curve  $C: \vec{r} = \cos t \vec{i} + \sin t \vec{j} + 3t \vec{k}$  where  $0 \le t \le 2\pi$ .
- 30. Test the continuity of f(x, y) defined by

$$f(x, y) = \frac{xy}{x^2 + y^2}$$
,  $(x, y) \neq (0, 0)$  and  $f(x, y) = 0$ ,  $(x, y) = (0, 0)$ .

- 31. Find the equation to the tangent plane and normal line to the surface  $f(x, y, z) = x^2 + y^2 + z^2 9 = 0$  at the point (1, 2, 4).
- 32. Evaluate the area enclosed by the Lemniscate  $r^2 = 4 \cos 2\theta$  using double integrals.

Turn over

- 33. Find the Local extreme values of  $f(x, y) = 3y^2 2y^3 3x^2 + 6xy$ .
- 34. Evaluate the volume of the region bounded by  $x^2 + y^2 = 4$ , y + z = 3, z = 0.
- 35. Show that  $\vec{f} = y \sin z \vec{i} + x \sin z \vec{j} + xy \cos z \vec{h}$  is conservative and find its scalar potential.

 $(6 \times 7 = 42 \text{ marks})$ 

### Part D

Answer any two questions.

Each question carries 13 marks.

- 36. (a) State Gauss divergence theorem and use it to evaluate the outward flux of  $\vec{f} = xy \vec{i} + yz \vec{j} + xz \vec{k}$  through the surface of the cube cut from the first octant by the planes x = y = z = 1.
  - (b) If S is a closed surface enclosing a volume V, then prove that  $\int \int_{S} \vec{r} \cdot n dS = 3V$ .
- 37. (a) Evaluate the surface integral  $\int_{\mathbb{R}} \vec{f} \cdot ndS$  where  $\vec{f} = y \vec{i} + x \vec{j} + z^2 \vec{k}$  over the cylindrical surface S given by  $x^2 + y^2 = a^2$ , z = 0, z = h.
  - (b) Find angle between the surfaces  $x^2 + y^2 + z^2 = 9$  and  $z = x^2 + y^2 3$  at (2, -2, 2).
- 38. (a) Find the value of  $\int_{(1,0,0)}^{(0,1,0)} \sin y \cos x dx + \cos y \sin x dy + dz$ 
  - (b) In what direction from the point (2, 1, -1) the directional derivative of  $\phi(x, y, z) = x^2 yz^3$  is maximum and find the magnitude of this maximum.

 $(2 \times 13 = 26 \text{ marks})$